

This IssueViews **99,269** | Citations **34** | Altmetric **1045** | Comments **2**

PDF



CME & MOC



Share



Comment

Editorial**FREE**

May 20, 2020

Nasal ACE2 Levels and COVID-19 in ChildrenAnkit B. Patel, MD, PhD¹; Ashish Verma, MBBS¹[» Author Affiliations](#) | [Article Information](#)

JAMA. 2020;323(23):2386-2387. doi:10.1001/jama.2020.8946

**COVID-19 Resource Center**Related
Articles

Coronavirus disease 2019 (COVID-19) has disproportionately affected certain vulnerable populations. Studies noted higher rates of certain comorbidities such as hypertension, diabetes mellitus, and chronic obstructive pulmonary disease in patients infected with COVID-19 with severe disease.¹ Additionally, areas with more racial/ethnic minorities and higher rates of poverty have been shown to have higher rates of COVID-19 hospitalization and death.² After adjustment for comorbidities, age has been independently associated with increased mortality due to COVID-19.³ However, limited attention has been given to children, who appear to have lower risk of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and mortality.

In this issue of *JAMA*, Bunyavanich et al⁴ identify a possible factor that may be related to lower rates of SARS-CoV-2 infection in children. The authors evaluated gene expression in nasal epithelial samples collected as part of a study involving patients with asthma from 2015 to 2018. The nasal epithelium is one of the first sites of infection with SARS-CoV-2, and the investigators probed for the expression of the cell surface enzyme angiotensin-converting enzyme 2 (ACE2), which has been proven to bind to SARS-CoV-2 spike protein and promote internalization of the virus into human cells.⁵ Among a cohort of 305 patients aged 4 to 60 years, older children (10-17 years old; n=185), young adults (18-24 years old; n=46), and adults

PDF

Help

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

children (4-9 years old; n=45), and *ACE2* expression was higher with each subsequent age group after adjusting for sex and asthma.

Numerous studies have highlighted the low rates of SARS-CoV-2 infection in children compared with adults. Children have been shown to have fewer and less severe symptoms compared with adults.^{6,7} This leads to the question of whether low rates of SARS-CoV-2 infection in children are due to low rates of testing in children, or if children are less susceptible to infection. An evaluation of 1286 close contacts of index cases in China found that infection rates in children were comparable with or slightly higher than in younger adults (aged 30-49 years) but were significantly lower than in older patients (aged ≥ 60 years).⁸ This finding suggests that children seem to have similar rates of becoming infected compared with middle-aged adults following close contact with a person infected with SARS-CoV-2. In contrast, a targeted screening approach in Iceland found SARS-CoV-2 in 6.7% of children younger than 10 years old (n=564) compared with in 13.7% of people aged 10 years or older (n=8635). A population-wide screening approach not restricted to people at risk of disease in the same study found no positive cases of SARS-CoV-2 among children under 10 years old, whereas 0.8% of people older than 10 years were positive for SARS-CoV-2.⁹ Although the true rate of acquiring SARS-CoV-2 infection among children of different age groups in the United States remains unclear, initial studies have suggested a lower incidence of SARS-CoV-2 infection in US children.⁷

While ACE2 binds to the receptor binding domain of SARS-CoV-2, a serine protease, TMPRSS2, is shown to cleave SARS-CoV-2 S protein, allowing for cell membrane fusion and endocytosis of the virus and subsequent viral replication. To better understand expression of these 2 key proteins, a number of single-cell RNA sequencing experiments have mapped them to various human cell types. One study using human and nonhuman primate tissue found *ACE2* expression in the secretory cells of the nasal epithelia (upper respiratory tract) and in the type II pneumocytes of the lower respiratory tract.¹⁰ In fibrotic human lung tissue, only 1.4% of type II pneumocytes expressed *ACE2*, and 0.8% of type II pneumocytes expressed both *ACE2* and *TMPRSS2*. In human ethmoid sinus surgical specimens representing upper respiratory epithelia, 1.3% of all secretory cells expressed *ACE2* while 0.3% expressed both *ACE2* and *TMPRSS2* but were further enriched in secretory goblet cells.¹¹ The spatial resolution offered by single-cell RNA sequencing highlights the limited number of cells in the respiratory tract that express *ACE2*. Further evaluation of *ACE2* expression at the single-cell level in children could help explain whether a lower percentage of *ACE2*-expressing cells or a decrease in expression in *ACE2* expression per cell is related to lower *ACE2* expression in nasal epithelium, as noted in children identified by Bunyavanich et al.⁴

ACE2 has an important role in counterbalancing the effects of ACE. Angiotensin II, a product of ACE cleaving angiotensin I, can cause vasoconstriction, inflammation, and fibrosis by signaling through angiotensin II type 1 receptors. *ACE2* can cleave angiotensin II to angiotensin 1-7, which can suppress inflammation and fibrosis and generate vasodilation by binding to the Mas receptor. Previous studies have found *ACE2* to

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

serves as a receptor for viral entry, then is more *ACE2* or less *ACE2* expression protective for children? In the nasal epithelium of the upper airway, lower *ACE2* expression could be helpful in decreasing acquisition of SARS-CoV-2 infection. However, in the lower respiratory tract, it appears that decreased *ACE2* expression could portend a higher risk of developing severe acute respiratory distress and lung injury.

Bunyavanich et al⁴ suggested that *ACE2* expression in the nasal epithelium in their cohort does not reflect *ACE2* expression in the pulmonary epithelium, and the expression of *ACE2* in the lower respiratory tract is under different regulation. This emphasizes the importance of understanding the distribution of *ACE2* in cells in different parts of the respiratory epithelium but also between cell-bound and plasma fractions. *ACE2* is cleaved from the cell membrane on binding with SARS-CoV-2, releasing *ACE2* into the plasma. The role of soluble *ACE2* in neutralizing SARS-CoV-2 virus has recently been shown in vitro,¹³ but its activity in vivo is still to be determined. Ultimately, studying tissue expression of *ACE2* in the lower respiratory tract of children may be helpful in understanding differences in the severity of COVID-19 among children compared with adults.

A new study called Human Epidemiology and Response to SARS-CoV-2 (HEROS), funded by the National Institute of Allergy and Infectious Diseases, is designed to prospectively follow 6000 children to determine risk factors for development of COVID-19.¹⁴ Data from this study could help identify whether the lower *ACE2* expression identified by Bunyavanich et al⁴ correlates with lower rates of SARS-CoV-2 infection, and could serve to support the possibility that decreasing *ACE2* expression in the nasal epithelium may be a potential therapeutic approach to mitigate transmission of COVID-19.

Advertisement

Article Information

[Back to top](#)

Corresponding Author: Ankit B. Patel, MD, PhD, Brigham and Women's Hospital, 75 Francis St, Boston, MA 02132 (apatel38@bwh.harvard.edu).

Published Online: May 20, 2020. doi:[10.1001/jama.2020.8946](https://doi.org/10.1001/jama.2020.8946)

PDF

Help

Conflict of Interest Disclosures: Dr Patel reported receipt of grants from Relypsa and consulting fees from Third Rock Ventures. No other disclosures were reported.

References

1. Richardson S, Hirsch JS, Narasimhan M, et al; Northwell COVID-19 Research Consortium. Present-

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

- in the New York City area. *JAMA*. Published online April 22, 2020. doi:[10.1001/jama.2020.6775](https://doi.org/10.1001/jama.2020.6775)
[Article](#) | [PubMed](#) | [Google Scholar](#)
2. Wadhera RK, Wadhera P, Gaba P, et al. Variation in COVID-19 hospitalizations and deaths across New York City boroughs. *JAMA*. Published online April 29, 2020. doi:[10.1001/jama.2020.7197](https://doi.org/10.1001/jama.2020.7197)
[Article](#) | [PubMed](#) | [Google Scholar](#)
 3. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in COVID-19. *N Engl J Med*. Published online May 1, 2020. doi:[10.1056/NEJMoa2007621](https://doi.org/10.1056/NEJMoa2007621)
[PubMed](#) | [Google Scholar](#)
 4. Bunyavanich S, Do A, Vicencio A. Nasal gene expression of angiotensin-converting enzyme 2 in children and adults. *JAMA*. Published online May 20, 2020. doi:[10.1001/jama.2020.8707](https://doi.org/10.1001/jama.2020.8707)
[Article](#) | [Google Scholar](#)
 5. Zhou P, Yang X-L, Wang X-G, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273. doi:[10.1038/s41586-020-2012-7](https://doi.org/10.1038/s41586-020-2012-7)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
 6. Castagnoli R, Votto M, Licari A, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review. *JAMA Pediatr*. 2020;174(9). doi:[10.1001/jamapediatrics.2020.1467](https://doi.org/10.1001/jamapediatrics.2020.1467)
[Article](#) | [PubMed](#) | [Google Scholar](#)
 7. Bialek S, Gierke R, et al; CDC COVID-19 Response Team. Coronavirus disease 2019 in children—United States, February 12–April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):422-426. doi:[10.15585/mmwr.mm6914e4](https://doi.org/10.15585/mmwr.mm6914e4)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
 8. Bi Q, Wu Y, Mei S, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis*. 2020;S1473-3099(20)30287-5. doi:[10.1016/s1473-3099\(20\)30287-5](https://doi.org/10.1016/s1473-3099(20)30287-5)
[PubMed](#) | [Google Scholar](#)
 9. Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS-CoV-2 in the Icelandic population. *N Engl J Med*. Published online April 14, 2020. doi:[10.1056/NEJMoa2006100](https://doi.org/10.1056/NEJMoa2006100)
[PubMed](#) | [Google Scholar](#)
 10. Sungnak W, Huang N, Bécavin C, et al; HCA Lung Biological Network. SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med*. 2020;26(5):681-687. doi:[10.1038/s41591-020-0868-6](https://doi.org/10.1038/s41591-020-0868-6)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)

PDF

Help

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

11. Ziegler CGK, Allon SJ, Nyquist SK, et al. SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues. *Cell*. Published online April 27, 2020. doi:[10.1016/j.cell.2020.04.035](https://doi.org/10.1016/j.cell.2020.04.035)
[Google Scholar](#)
12. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*. 2005;436(7047):112-116. doi:[10.1038/nature03712](https://doi.org/10.1038/nature03712)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
13. Monteil V, Kwon H, Prado P, et al. Inhibition of SARS-CoV-2 infections in engineered human tissues using clinical-grade soluble human ACE2. *Cell*. 2020;S0092-8674(20)30399-8. doi:[10.1016/j.cell.2020.04.004](https://doi.org/10.1016/j.cell.2020.04.004)
[PubMed](#) | [Google Scholar](#)
14. Study to determine incidence of novel coronavirus infection in US children begins. National Institutes of Health website. Published May 4, 2020. Accessed May 11, 2020.
<https://www.nih.gov/news-events/news-releases/study-determine-incidence-novel-coronavirus-infection-us-children-begins>

Comment

2 Comments for this article

EXPAND ALL

May 23, 2020

Gene Expression vs Protein, and Asthma

Dora Smith, None | None

PDF

Help

1. The original article mentioned that ACE2 gene expression is not the same thing as ACE receptors and they may not be correlated - but the editorial does not.

2. The original article mentioned that 45% of the children in the study had asthma, and that having asthma was controlled for. However, the article does not report what relationship existed between ACE2 gene expression and asthma, and they surely noticed if there was such a relationship. This matters because people with asthma have lower ACE2 levels - gene expression or receptors - and are also, worldwide, ...

READ MORE

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

May 26, 2020

Nasal ACE2 Levels and Fructose Consumption as a Risk Factor For Covid19

Declan Waugh, BSc Env Science | Research Scientist

I read the research letter by Bunyavanich et al. and accompanying editorial in JAMA with interest. The authors identified that ACE2 gene expression was significantly higher in older children, young adults and adults and suggested that lower ACE2 expression in children may be a possible factor that may be related to lower rates of SARS-CoV-2 infection in children. While the molecular mechanisms that mediate ACE2 expression are poorly understood, it has been observed that fructose metabolism increases ACE2 expression [1]. Moreover, fructose consumption is also directly implicated in the pathogenesis of hypertension, obesity, diabetes, kidney disease, cardiovascular disease, and ...

[READ MORE](#)

See More About

- Infectious Diseases
- Otolaryngology
- Pulmonary Medicine
- Coronavirus (COVID-19)
- Pediatrics



[Coronavirus Resource Center](#)

SCHEDULED MAINTENANCE

Our websites may be periodically unavailable between 7:00PM CT July 10, 2021 and 1:00AM CT July 11, 2021 for regularly scheduled maintenance.

[PDF](#)

[Help](#)

Trending

Research

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

July 6, 2021

Research 

Effect of 2 Inactivated SARS-CoV-2 Vaccines on Symptomatic COVID-19 Infection in Adults

July 6, 2021

Research 

Association Between BNT162b2 Vaccination and SARS-CoV-2 Infection Incidence Among Health Care Workers

June 22, 2021

Select Your Interests

Advertisement

JOB LISTINGS ON JAMA CAREER CENTER®

CLINICAL FACULTY: CARDIOVASCULAR INTENSIVIST

Tucson, Arizona

Physician (Critical Care)

Ft. Hood, TX

CARDIAC INTENSIVIST

Phoenix, Arizona

Physician, Critical Care

Oklahoma City, OK

Physician-Critical Care-St. Agnes Hospital

Fond du Lac, WI

See more at JAMA Career Center

PDF

Help

Others Also Liked

COVID-19 and the gastrointestinal tract: more than meets the eye

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)

ACE2 expression and sex disparity in COVID-19

Maria Cristina Gagliardi et al., Cell Death Discov, 2020

Kidney injury molecule-1: a novel entry factor for SARS-CoV-2

Cheng Wan, Journal of Molecular Cell Biology, 2021

Powered by **TREND MD**

Trending

COVID-19 Seroprevalence and Risk Factors Among Health Care Workers in Italy

JAMA Network Open | *Research* | July 6, 2021

Serologic Status and Toxic Effects of the BNT162b2 Vaccine in Patients With Cancer

JAMA Oncology | *Research* | July 8, 2021

Effect of Bamlanivimab vs Placebo on Incidence of COVID-19 at Skilled Nursing and Assisted Living Facilities

JAMA | *Research* | July 6, 2021

PDF

Help

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [Cookie Policy](#) | [Continue](#)